



CAMET Regional Workshop on Metrology and Technology Challenges of Climate Science and Renewable Energy Guatemala, Guatemala

Gaatomala, Gaatomala

Jorge Koelliker D., Dr. rer. nat., CENAM

Air Quality Monitoring and Climate Change Related Measurements (GHG)

CONTENT

- 1. Climate change issues and consequences
- 2. Quantities to be measured?
- 3. Air quality and GHG substances
- 4. Regulatory framework
- 5. Reflections and conclusions





1. Climate change issues and consequences

- Increase of the average temperature in the biosphere due to the increase of more heat absorbing molecules in the air and the natural greenhouse effect
 - Increase in frequency, intensity and change of pattern in meteorological phenomena, e. g.: rain, hurricanes, storms, more prolonged droughts and heating waves.
 - Glaciar depletion or extinction of ice in the poles and in the terrestrial surface in the mountains
 - Increase in temperature of see and in its level
 - Change in the behaviour of animals and plants

More consequences

 Loss of land in coasts; floods; habitat, species and agricultural losses; increase of some bacteria and parasites illnesses associated with heat; ...





2. Quantities to be measured?

Recognize quantities not just variables

- Meteorological quantities: temperature, wind speed and direction, water vapour amount fraction, pressure, precipitation, surface/earth radiation budget, solar irradiance, cloud properties, ...
- Atmospheric composition: long-lived GHGs: amount fractions of CO₂, CH₄, N₂O, ...; O₃, aerosol and their precursors
- Oceans: CO₂ partial pressure, O₂, acidity, phytoplanckton, temperature,

•





The Greenhouse gases: Half live time and warming potential

(Origen	Gases	Fuentes		Potencial de Calentamiento
	ases de origen natu	Bióxido de carbono (CO₂)	Quema de combustibles fósiles (carbón, derivados de petróleo y gas), reacciones químicas en procesos de manufactura; (como la producción de cemento y acero) cambio de uso de suelo (deforestación).	50 a 200	1
		Metano (CH4)	Descomposición anaerobia (cultivo de arroz, rellenos sanitarios, estiércol), escape de gas en minas y pozos petroleros.	12 ± 3	21
		Óxido nitroso (N₂O)	Producción y uso de fertilizantes nitrogenados, quema de combustibles fósiles.	120	310
	ıses antropogénico	Hidrofluorocarbonos (HFCs)	Emitidos en procesos de manufactura y usados como refrigerantes.	1.5 a 264	140-11,700
		Perfluorocarbonos (PFCs)	Producción de Aluminio, fabricación de semiconductores, sustituto de las sustancias destructoras del ozono. Ej. Uso de solventes, espumas, refrigeración fija.	2600 a 50000	6,500-9,200
		Hexafluoruro de Azufre (SF ₆)	Producción y uso en equipos eléctricos; Producción de magnesio y aluminio; Fabricación de semiconductores.	3200	23,900





Why Greenhouse gases and air pollutants together?

"A gaseous atmosphere is maintaining our life"

- <u>Significant</u>: understanding chemistry and physics of the atmosphere
 - Primary and secondary air pollutants and their precursors or (sub)products are sometimes all relevant pollutants in climate change (e. g. O₃: CH₄, VOCs, NOx, etc.; CO₂: CO, CH₄, carbon cycle; nitrogen cycle; acid rain: SOx, NOx)...
 - Substances and radicals that induce reactions or the presence of other air pollutants (CHCO, OH, PANs, etc.). Some of them are inestable, thus not prone to have them as measurement standards.





- Significant air pollutants in atmospheric monitoring (emissions and inmissions)

 Development of Chemical measurement standards in Mexico
 - Greenhouse gases (GHGs): those 7 reported groups GHGS to UNFCCC (& Kyoto Protocol)
 - CO₂, CH₄, N₂O, SF₆, originally chlorofluorocarbons (CFCs), replaced by hydrochlorofluorocarbons (HCFCs), now replaced by the latest hydrofluorocarbons (HFCs), NF₃
 - Short-Lived Climate Forcers or Pollutants (SLCFs / SLCPs) for near-term climate protection: CH₄,
 tropospheric O₃, black carbon, short-lived hydroflourocarbons (HFCs).
 - HFCs are a type of fluorinated greenhouse gas intentionally made as replacements for stratospheric ozone depleting substances (ODS, *Montreal Protocol*), for use in the same applications (air conditioning, refrigeration, solvents, foam blowing and aerosols)
 - Criteria air pollutans (gases) for ambient air (inmissions): are 7, but in Mexico only testing 5
 - Pb, CO, O₃, NO₂, PM_{10, 2,5 ... 1.0 ...,} SO₂, VOCs, H₂S (not a criteria air pollutant, but at ambient levels also measured in specific places)
 - Stack and mobile emissions
 - Metals (Pb, Hg, ...), hydrocarbons (HCs expressed as C₃H₈ or bencene) CO, CO₂, NO_x, PM_{10, 2.5, ..., 1,0, ...} SO₂, VOCs, ...
 - Persistant Organic Pollutants (POPs). Stockholm Convention. Dioxins, Pesticides, PCBs.
 - PAHs (carcinogenic mutagenic).







What has been already done and is being done? Comparisons in Air quality & GHG-CCQM-CIPM


Date updated	1 Dec 2013					
WG	Reference No.	Description	Coordinating Laboratory	Start date	Status	Hyperlink to latest version of report
Criteria air polluta	ants			_		
GAVG	BIPM.QM-K1	Ozone at ambient level	BIPM	2006	Approved for Equivalence	BIPM QM K1 Reports
GAVG	CCQM-K51	CO in nitrogen (5 µmol/mol)	NMISA	2006	Approved for Equivalence	CCQM-K51 Final Report
GAVG	CCQM-K84	Ambient Carbon Monoxide	KRISS	2010	In progress	
GAVG	CCQM-K74	Nitrogen dioxide 10 μmol/mol	BIPM	2010	Approved for Equivalence	CCQM-K74 Final Report
GAVG	CCQM-P73	Nitrogen monoxide in nitrogen	BIPM	2006	Completed	CCQM-P73 Final Report
GAVG	CCQM-K26.a	Reactive gases-ambient levels - NO in N₂	NPL	2003	Approved for Equivalence	CCOM-K26.a_Final_Report
GAVG	CCQM-K26.b	Reactive gases-ambient levels - SO2 in Air	NPL	2003	Approved for Equivalence	CCGM-K26.b_Final_Report
Other air pollutan	ts		02			
GAVG	CCQM-K7	Benzene/toluene/xylene (BTX) in N₂/Air	NIST S	1999	Approved for Equivalence	CCOM-K7_Final_Report
GAWG	CCQM-K10	BTX in N2 (low conc 10-30 ppb)	NIST/NPL	2001	Approved for Equivalence	CCOM-K10_Final_Report
GAVG	CCQM-K22	VOCs in Air	NMIJ	2003	Approved for Equivalence	CCOM-K22_Final_Report
GAWG	CCQM-K121	Terpenes in air	NIST	2016	Planned	
GAVG	CCQM-K46	Ammonia in nitrogen	VSL	2005	Approved for Equivalence	CCOM-K46_Final_Report
GAWG	CCQM-K117	NH₃ in nitrogen	VSL	2015	Planned	
GAVG	CCQM-K90	Formaldehyde in air	вірм	2010	Planned	
Climate change (GHG)					
GAVG	CCQM-P41	Greenhouse gases CO ₂ , CH ₄ - ambient levels	VSL	2002	Completed	CCOM-P41_Final_Report
GAVG	CCQM-K52	CO₂in air (360 - 400 ppm)	VSL	2006	Approved for Equivalence	CCQM-K52 Final Report
GAWG	CCQM-K120	Ambient CO ₂	BIPM with NIST	2015	Planned	
GAVG	CCQM-K82	Methane in air	BIPM	2009	In progress	
GAVG	CCQM-K68	N2O at ambient levels	KRISS	2008	Approved for Equivalence	CCQM-K68 Final Report
GAVG	CCQM-K15	SF ₆ , CFCs - emission levels	KRISS	2003	Approved for Equivalence	CCOM-K15_Final_Report
GAVG	CCQM-K94	DMS (Di-methyl sulphide) in nitrogen	KRISS	2011	In progress	
GAVG	CCQM-K83	Halocarbons in air	NIST	2011	In progress	





What has been already done and is being done? Comparisons in Air quality & GHG-CCQM-CIPM

Criteria air pollutants: Ozone








What has been already done and is being done? Comparisons in Air quality & GHG-CCQM-CIPM

Criteria air pollutants: Carbon monoxide, a new CCQM-K84 is in progress








What has been already done and is being done? Comparisons in Air quality & GHG-CCQM-CIPM

Criteria air pollutants: Nitrogen dioxide: 1st Nitrogen monoxide








What has been already done and is being done? Comparisons in Air quality & GHG-CCQM-CIPM

Criteria air pollutants: Nitrogen dioxide: 2nd Nitrogen dioxide








What has been already done and is being done? Comparisons in Air quality & GHG-CCQM-CIPM

Criteria air pollutants: Sulfur dioxide











What has been already done and is being done? Comparisons in Air quality & GHG-CCQM-CIPM

Other (criteria) air pollutants: some VOCs (only Hazarduos Air Pollutans: Benzene, chloroform, dichloromethane, trichloroethylene, tetrachloroethylene, 1,2-dichloroethane,1,3-butadiene, and vinyl chloride) at 100 nmol/mol level; BTEX 50 to 100 nmol/mol, e. g.














What has been already done and is being done? Comparisons in Air quality & GHG-CCQM-CIPM

Other air pollutants: Ammonia in N_2 , 30 μ mol/mol to 50 μ mol/mol








What has been already done and is being done? Comparisons in Air quality & GHG-CCQM-CIPM

CCQM-P41 Greenhouse gases – CO₂ and CH₄ in Air (2003)



Example of CRMs and DQO of WMO



Projected Typical Certified Values for SRM 1720 : Northern Hemisphere Greenhouse Gases

		NIST Concentration ^a	NOAA Concentrationa	WMO Data Quality Objective ^b
	CO ₂	393. 30 ± 0.10 μmol/mol	393.38 ± 0.20 μmol/mol	± 0.1 μmol/mol
	CH ₄	1878.1 ± 1.0 nmol/mol	1875.9 ± 1.5 nmol/mol	± 2 nmol/mol
	N_2O	323.04 ± 0.20 nmol/mol	323.10 ± 1.0 nmol/mol	± 0.1 nmol/mol
	СО	≈ 160 ± 4 nmol/mol (vary 35 % sample to sample)	No NOAA Data	± 2 nmol/mol
	SF ₆	≈ 7 ± 0.04 pmol/mol	7.16 ± 0.04 pmol/mol	± 0.02 pmol/mol

^a Expanded uncertainty at k=2 coverage factor, 95 % confidence interval.

^b Standard uncertainty at 1 σ (k=1 coverage factor).








What has been already done and is being done? Comparisons in Air quality & GHG-CCQM-CIPM

GHGs: Carbon dioxide, a new CCQM-K120 planned









What has been already done and is being done? Comparisons in Air quality & GHG-CCQM-CIPM

GHGs: Methane. Data quality objective (DQO) of the WMO for CH₄ almost achieved, but not yet in the uncertainty of gaseous PSMs. DQO of the WMO defined as metrological compatibility of measurement results (VIM, 2.47)










What has been already done and is being done? Comparisons in Air quality & GHG-CCQM-CIPM

GHGs: Nitrous oxide, 320 nmol/mol









What has been already done and is being done? Comparisons in Air quality & GHG-CCQM-CIPM


GHGs: Sulfur hexaflouride and CFCs at emission levels (Carbon tetraflouride) 100 μml/mol to cover CF₄, C₂F₆, CHF₃, SF₆, and NF₃



























Mitigation

- National policy of mitigation should include: diagnostic (national and sectorial baseline), plan, measurement, monitoring, report, verification and evaluation of national emissions.
- Arrangements at federal, state and municipal level to reduce emissions at specific sectors and activities through actions and programs.
- International treaties signed by Mexico to be considered.
- Low-emission development strategies (LEDS)
- Green Economy
- Nationally Appropriate Mitigation Activities (NAMAs)
- Mexico is a member of the Global Green Growth Institute since 2012
- Mexico is a founding member of Climate and Clean Air Coalition (CCAC) to Reduce Short-Lived Climate Pollutants

Policies

Climate science, <u>mitigation</u> and adaptation to CC

Actions (for Sustainable Development)


Technology



National Strategy on Climate Change Vision 10-20-40 (years) 2013

Without metrology


The measurement, report and verification (MRV) for mitigation of GHGs should be better supported by Metrology and quality infrastructures in each country. Cooperation of authorities with NMIs have to be closer for these tasks. However, I think that the MRV process is not yet at all considering Metrology, because it could be seen as an additional requirement to promote effective reductions. Same situation in NAMAs. So convince also IPCC national experts







Fifth National Communication to the UNFCCC (for 2010, reported 2012) - Mexico








Reflections

Case of 40 Megacities: 3rd contributor to GHGs after USA and China



Cities: responsible for 70% of fossil-fuel CO2 emissions

A 10-km-resolution map indicates the distribution and intensity of fossil fuel CO2 emission sources. The regions with greatest emission intensity are indicated by red and black (urbanized areas and associated large power plants). The black circles indicate a vision for future surface measurement networks concentrated within the 23 existing megacities. Blue circles indicate the 14 additional megacities projected to exist by 2025. The dashed rectangles indicate the fields of regard of three remote-sensing instruments that if hosted on geostationary satellites would offer sustained, wall-to-wall mapping of nearly every emission source. The satellite and surface network data, integrated with improved high-resolution emission estimates would provide a robust system for assessing and informing policies. Map: (EDCAR version 4.0) 2009.





Reflections

- CENAM has 16 years experience on emission services: vehicle and stack emissions, participation in some CCQM-Key Comparisons and Pilot Studies of GHGs and Air Pollutants. It is the only Iberoamerican NMI with chemical CMCs as core capabilities in gases (in the MRA-CIPM) published today (it does not include some GHGs, GHG chemical measurement standards are challenging, not core!). As member of the CCQM we have regular participation in the GAWG and we are following comparisons and knowlegde in GHGs and air quality measurements.
 - Steps of Development in Gas Metrology: please first binary gas mixtures high concentrations, i.e. emissions, lower concentration and multicomponent gas mixtures are challenging: so truly air quality levels (not their premixtures) are very challenging!, i. e. inmissions are more difficult!
- Politically, Mexico has been very involved in climate change initiatives. In my knowledge, Mexico is the 2nd country in the World with a General Law of Climate Change (2012, first was UK), having a new National Institute of Ecology and Climate Change. First country reporting to the UNFCCC inventories on Short Live Climate Forces (SLCF, 2012).





Reflections

- Many scientist doing an excellent job in monitoring the air, but only a few truly understand and apply Metrological Traceability and Metrology in Chemistry Capabilities, so more guidelines are required.
 - Not only for gas analysis in gas phase, but also for wet methods. Examples:
 - In spectroscopic measurements doing "absolute measurements" linestrengths have many times very high uncertainties and no traceability, there are significant uncertainty sources in emerging technologies, e. g. for QCLs, CRDS, etc.
 - Wet methods also need good metrological practices, e. g. measuring vertical profiles of O_3 in planes or globes absorbed in KI.
- So we should review the metrological quality of the stuties performed in our own country. Do not do only isolated efforts: synergy is needed!





Conclusions

- Better quality of weather and air quality data for forecast or/and complement/substitution to measurements with models. Work on metrics of models. What bias, if allowed, is okay between traceable measurements and models?
- Synergy: cooperate closer with Environmental Authorities and Universities (e.g. air pollution and models), but also with the National Meteorological Service (SMN) and Organizations (State or Municipal levels) in charge of Meteorological Stations.
- Development of traceable and reference methods for assays of GHGs.
- Set of governmental priorities is clear, but in Metrology very delayed: SLCF and particles: innovation opportunities in metrological traceability.
- Contribute with traceable results in inventory guidelines of the IPCC and MRV of mitigation actions and NAMAs and initiate/continue Metrology Support to the inventory reports for the UNFCCC.



